



Penetration of foliar-applied Zn and its impact on apple plant nutrition status

Ruohan Xie¹, Jianqi Zhao¹, Lingli Lu¹, Patrick Brown², Jiansheng Guo^{3, 4} and Shengke Tian^{1*}

¹College of Environmental & Resource Science, Zhejiang University, Hangzhou, China

²Department of Plant Sciences, University of California, Davis, CA, USA

³Department of Pathology of Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China.

⁴Center of Cryo-Electron Microscopy, Zhejiang University School of Medicine, Hangzhou, China.

INTRODUCTION

Vj g"cdutrvkqp"qh"hnkct"htvkl gt"ku"e"eqo rrgz"rtqegu"cpf"ewtgpv"tgo ckpu"rqqr"wpf gtuvqf O'Vj g"cdkkl"vq"xkwrkl g"cpf"s wcpvkh"vj g"rcvj y c{"vj cv"grgo gpvu"veng"hmvy kpi"cr rdekvp"vq"ngch"uwthegu"ku"etkklcn"ht"vj g"uelgpeg"cpf"r tcevecn"cr rdekvp"qh"hnkct"htvkl gt O'Cr r ng" **Malus domestica* Dqtnj 0'y cu'uggevfg"cu"cp"gzr gkto gpvcl'ur gelgu"dgecwug"kv"ku"j ki j n"ugpukxg"vq"\ p"f ghelgpe{ O' Oquv"cr r ng"qtej ctf u"kp"ctgeu"y kj"ecrectgqu"cpf"ucn/chhgevfg"uqku"uwthgt"htqo "\ p"f ghelgpe{"gxgt{"{gct."y j lej"tguwmu"kp"j ki j"nqu"qh"lgrf"cpf"ugxgtg"f gvgtktcvqp"qh"htwk"s wcrkl{

MATERIALS & METHODS

D{"wulpi"o ketq"Z/tc{"hqtguepeg."y g"ctg"cdng"vq"cpcl{| g"vj g"ko rcev"qh"uwthceg"r j {uleqej go kcln"rtqr gtvku"qh"cr r ng"r rcpv"ngcxgu"qp"vj gkt"\ p"cdutrvkqp"cpf"htvj gt"vcpurqecvqp O'Vj gp."j ki j gt"tguqnwkqp"ko ci kpi"y cu"cr r rkgf"vq"cuugu"vj g"r gpgvcevkqp"rtqegu"d{"kp"xkq"o cr r kpi"qh"\ p"cvuwdegmwt"rgxgr O'Y g"cuq"kp xguvki cvg"vj g"kpvtcevkqp"qh"\ p"cpf"qvj gt"grgo gpvu"vq"gzco kpg"j qy"fq"pwtklqpcn"ucwu"kp"cr r ng"ngcxgu"tgr qp"vq"hnkct"\ p"cr r dekvp O'Vj g"cdkkl"vq"xkwrkl g"cpf"s wcpvkh"vj g"rcvj y c{"vj cv"grgo gpvu"veng"hmvy kpi"cr r dekvp"vq"ngch"uwthegu"ku"etkklcn"ht"qwt"wpf gtuvcpf kpi"qh"vj g"wkkl"qh"hnkct"htvkl gtu"cpf"cmvy u"ht"vj g"fkgev"eqpvcu"qh"tgrvkg"ghhece{"qh"f hhtgpv"htto wrcvqpucpf"ej go kwtkgu O

RESULTS & CONCLUSIONS




Figure 1 SEM micrographs of adaxial apple leaves (A) and Micro-XRF mapping of Zn in cross-sections of leaves after foliar Zn application (B)




Figure 2 SEM micrographs of abaxial apple leaves with high, medium and low trichome densities (A) and Micro-XRF mapping of Zn in cross-sections of leaves after foliar Zn application (B)






Figure 5 Zn levels in the leaves

Conclusion

Figure 3 Subcellular distribution of Zn in apple leaves after foliar Zn application. Nano-XRF images were obtained from cross sections of apple leaf tissue after foliar Zn treatment of adaxial (A) and abaxial (B) leaf surfaces.

The results indicate that the absorption of foliar applied Zn was largely dependent on plant leaf surface characteristics. Higher-resolution elemental maps revealed that high-binding capacity of cell wall for Zn contribute to the observed limitation of Zn penetration across epidermal cells. Trichome density and stomatal aperture had opposite effects on Zn fertilizer penetration; a higher density of trichomes increased the hydrophobicity of leaves, whereas the presence of stomata facilitated foliar Zn penetration. Low levels of Zn promoted the accumulation of other mineral elements in treated leaves, the complexation of Zn with phytic acid occurred potentially owing to the exposure to high Zn conditions. The present study provides the direct visual evidence for Zn penetration process across the leaf surface which is important for the development of strategies for Zn biofortification in crop species.